


One day.
One dose.
That's it!



solosec
secnidazole
2g Oral Granules
LUPIN

One **1** & Done



SOLOSEC[®] is a single dose oral antibiotic that provides a complete course of therapy to treat trichomoniasis (trich)¹

Inside you'll find helpful information, including:

- What is trich?
- What causes trich?
- Reasons why you need to see your healthcare provider
- Why treatment with SOLOSEC[®] is different

What is SOLOSEC[®]?

SOLOSEC[®] (secnidazole) 2g oral granules is a prescription medicine used to treat Trichomoniasis, a common sexually transmitted infection (STI), in people 12 years of age and older. People taking SOLOSEC[®] for Trichomoniasis should avoid having sex until they and their sex partners are treated with the same dose and at the same time, to prevent reinfection. It is not known if SOLOSEC[®] is safe and effective in children under 12 years of age.

Select Important Safety Information

- You should not use SOLOSEC[®] if you've had an allergic reaction to secnidazole, other ingredients of the formulation, or other nitroimidazole derivatives or have Cockayne syndrome


Please see complete Important Safety Information on page 11 of this brochure.

What is trichomoniasis (trich)?



Trich is a common sexually transmitted infection (STI) that can affect both women and men²

How did I get trich?



Trich is passed from partner to partner during sexual intercourse²

Trich is...

The most common nonviral STI in the world²

Caused by a parasite called *Trichomonas vaginalis*²

Found in both men and women, although it's more common in women²



If you have sex—especially unprotected sex—with someone who has trich, you can be infected²



There's no test that can tell you who originally gave you the infection



Since trich can last for months or even years without treatment, a trich diagnosis does not necessarily mean you were recently infected²



Unless you and your partner(s) are all treated at the same time, you are likely to be reinfected if you have sex²

Symptoms of trich in women include²:

- Frothy vaginal discharge that's yellowish or greenish
- Vaginal itching and irritation
- An unpleasant odor

Many people with trich (around 70%) do not have any symptoms²



Why do I need to see my healthcare provider (HCP) right away to treat trich?



Leaving trich untreated can have **serious consequences**, so it's important to see your HCP for effective treatment²

Research has shown that untreated or improperly treated trich is associated with increased risk of:

- Developing pelvic inflammatory disease (PID), which can threaten your fertility^{2,3}
- Complications with your pregnancy, including late term miscarriage, premature birth, and low birth weight²
- Developing cervical cancer, which can be life-threatening⁴
- Getting sexually transmitted infections (STIs) like herpes, chlamydia, gonorrhea, and HIV, and transmitting STIs to your partner²
- Complications with gynecological surgery²

✓ You can take action to help reduce the risk of reinfection²

Not completing your full course of treatment is associated with trich that sticks around⁵

If your partner isn't treated, you're likely to get trich again²

Protect yourself and your partner(s) by getting treatment from your HCP, like **single-dose SOLOSEC®**

Select Important Safety Information

- Vaginal yeast infections may occur with SOLOSEC® and require an antifungal treatment.

Please see complete Important Safety Information on page 11 of this brochure.

What if I have trich and bacterial vaginosis (BV) at the same time?




SOLOSEC® is the **only single dose oral antibiotic** that's FDA-approved to treat **both trich and BV¹**

Many women have trich and BV—another vaginal infection—at the same time⁶

In fact, studies show that around 70% of women with trich are also infected with the bacteria that cause BV⁶

How BV & Trich Coexist



Having BV and trich increases your risk of getting other STIs like herpes, gonorrhea, and chlamydia⁷

Select Important Safety Information

- The most common side effects of SOLOSEC® include headache, nausea, vomiting, diarrhea, abdominal pain, vaginal itching and a bad, bitter or metallic taste in your mouth (dysgeusia).

Please see complete Important Safety Information on page 11 of this brochure.

What about my partner?



With SOLOSEC[®], you and your partner(s) can complete treatment for trich with **just one oral dose** each at the same time¹

What is SOLOSEC?

SOLOSEC[®] is a **single dose oral antibiotic** that's FDA-approved to treat **trichomoniasis (trich)** in adults¹



- All partners should be treated with the same dose and at the same time to prevent reinfection^{1,2}
- After you're diagnosed with trich, make sure you refrain from sexual intercourse with your partner(s) until everyone is treated and your symptoms go away²

- Let your partner know that SOLOSEC[®] provides a complete course of treatment with one oral dose¹



Ask your HCP if they can provide you with a SOLOSEC prescription for your partner

SOLOSEC

- ✓ Is an effective antibiotic medication that treats trich with **just one oral dose**¹
- ✓ Is rapidly absorbed by your body and **goes to work quickly** to fight trich⁸
- ✓ Keeps working in the body and treating trich **for several days after one dose**⁸
- ✓ **Was shown to be effective** in clinical studies – **92.3% of women** and **92%–100% of men** were cured after treatment with SOLOSEC^{®1,9}
- ✓ Isn't a pill. It's **different**¹



One day.
One dose.
That's it!



Select Important Safety Information

- Take SOLOSEC[®] exactly as your healthcare provider tells you to. Do not take SOLOSEC[®] more often than it is prescribed.

Please see complete Important Safety Information on page 11 of this brochure.

Select Important Safety Information

- You should not use SOLOSEC[®] if you've had an allergic reaction to secnidazole, other ingredients of the formulation, or other nitroimidazole derivatives or have Cockayne syndrome

Please see complete Important Safety Information on page 11 of this brochure.

Why is
SOLOSEC[®]
different?



One day.
One dose.
That's it!



SOLOSEC[®] lets you complete treatment with **one oral dose, one time**—not several pills over multiple days^{1,13,14}

How do I take
SOLOSEC[®]?



SOLOSEC[®] is easy to take¹



SOLOSEC[®] does not interfere with a common birth control pill*, condoms or diaphragms¹

*Combination ethinyl estradiol and norethindrone



With SOLOSEC[®], you can continue to use your vaginal products (tampons)

Complete Treatment with One Oral Dose

SOLOSEC[®] 2 g
1 Dose, ONCE

Another common medication for BV
14 Doses, 2x per day for 7 days



1 Open the packet and sprinkle the entire contents of the packet onto applesauce, yogurt or pudding



2 Stir and mix



3 Consume all of the mixture within 30 minutes. **DO NOT CHEW OR CRUNCH THE GRANULES**



You can drink a glass of water after taking SOLOSEC[®] to help with swallowing, if needed. Do not try to dissolve the medication in water or any other liquid

✓ Your sexual partner(s) should be treated at the same time¹

Select Important Safety Information

- Vaginal yeast infections may occur with SOLOSEC[®] and require an antifungal treatment.

Please see complete Important Safety Information on page 11 of this brochure.

Select Important Safety Information

- Take SOLOSEC[®] exactly as your healthcare provider tells you to. Do not take SOLOSEC[®] more often than it is prescribed.

Please see complete Important Safety Information on page 11 of this brochure.

How can I save money on my SOLOSEC® prescription?



There are multiple ways to access a co-pay card that may help you save money on your SOLOSEC® prescription*



If commercially insured, most eligible patients may pay **as little as \$25** for SOLOSEC®.* If you don't have insurance or are paying cash, you may still pay as little as \$75

- Visit solosec.com/savings-card to text a card to your mobile device, get a card emailed to you, or download a card onto your computer
- Alternatively, get a physical card from your HCP
- Using your mobile device, scan the QR code on this brochure below



*Per fill. Not valid for any person eligible for reimbursement of prescriptions, in whole or in part, by any state- or federally funded program, including Medicare, Medicaid, or Medicaid, or where prohibited, taxed or otherwise restricted by law. Cash-pay patients may pay as little as \$75. Eligibility requirements and terms and conditions apply. See Terms and Conditions at solosec.com for full details.

What is SOLOSEC®?

SOLOSEC® (secnidazole) 2g oral granules is a prescription medicine used to treat bacterial vaginosis in females 12 years of age and older. SOLOSEC® is used to treat Trichomoniasis, a common sexually transmitted infection (STI), in people 12 years of age and older. People taking SOLOSEC® for Trichomoniasis should avoid having sex until they and their sex partners are treated with the same dose and at the same time, to prevent reinfection. It is not known if SOLOSEC® is safe and effective in children under 12 years of age.

How should I take SOLOSEC®?

SOLOSEC® is a single-dose therapy for oral use. Use SOLOSEC® by sprinkling an entire packet of SOLOSEC® onto applesauce, yogurt, or pudding. The entire dose should be taken at once, and finished within 30 minutes. Avoid chewing or crunching the granules. SOLOSEC® should not be taken by dissolving the granules in any liquid. Avoid drinking alcohol during treatment with SOLOSEC® and for 2 days (48 hours) after you take SOLOSEC®.

Important Safety Information

- You should not use SOLOSEC® if you've had an allergic reaction to secnidazole, other ingredients of the formulation, or other nitroimidazole derivatives or have Cockayne syndrome.
- Before taking SOLOSEC®, tell your healthcare provider about all of your medical conditions, including if you
 - Are pregnant or plan to become pregnant.
 - Are breastfeeding or plan to breastfeed. Breastfeeding is not recommended. You should not breastfeed for 96 hours (4 days) after taking SOLOSEC®.
- Vaginal yeast infections may occur with SOLOSEC® and require an antifungal treatment.
- The most common side effects of SOLOSEC® include headache, nausea, vomiting, diarrhea, abdominal pain, vaginal itching and a bad, bitter or metallic taste in your mouth (dysgeusia).
- Take SOLOSEC® exactly as your healthcare provider tells you to. Do not take SOLOSEC® more often than it is prescribed.

Call your doctor for medical advice on side effects. You are encouraged to report suspected side effects of prescription drugs to FDA at www.fda.gov/MedWatch also at 1-800-FDA-1088 or contact Lupin Pharmaceuticals, Inc. at 1-844-SOLOSEC (1-844-765-6732).

Please see full prescribing information for SOLOSEC®, available at www.solosec.com.

References: 1. SOLOSEC® [prescribing information]. Baltimore, MD: Lupin Pharmaceuticals, Inc; 2022. 2. Workowski KA, Bachmann LH, Chan PA, et al. CDC Sexually Transmitted Diseases Treatment Guidelines, 2021. *MMWR Recomm Rep* 2021;70(RR-04):1-192. 3. Wiringa AE, Ness RB, Darville T et al. *Trichomonas vaginalis*, endometritis and sequelae among women with clinically suspected pelvic inflammatory disease. *Sex Transm Infect* 2020;96:436-438. 4. Yang S, Zhao W, Wang H, et al. *Trichomonas vaginalis* infection-associated risk of cervical cancer: A meta-analysis. *Euro J Obstet Gynecol* 2018;228:166-173. 5. Kissinger P. *Trichomonas vaginalis*: a review of epidemiologic, clinical and treatment issues. *BMC Infect Dis* 2015;15(307):1-8. 6. Sobel JD, Subramanian C, Foxman B, Fairfax M, Gygax S. Mixed Vaginitis—More than Coinfection and with Therapeutic Implications. *Curr Infect Dis Rep* 2013;15:104-108. 7. Brotman RM. Vaginal microbiome and sexually transmitted infections: an epidemiologic perspective. *J Clin Invest*. 2011;121(12):4610-4617. 8. Nyirjesy P, Schwelbe JR. Secnidazole: Next-generation antimicrobial agent for bacterial vaginosis treatment. *Future Microbiol* 2018;13:507-524. 9. Muzny CA, Schwelbe JR, Nyirjesy P, et al. Efficacy and Safety of Single Oral Dosing of Secnidazole for Trichomoniasis in Women: Results of a Phase 3, Randomized, Double-Blind, Placebo-Controlled, Delayed-Treatment Study. *Clin Infect Dis* 2021.

One day.
One dose.
That's it!

solosec
secnidazole
2g Oral Granules
LUPIN

One **1** & Done

SOLOSEC® is a
single dose oral
antibiotic that
effectively
treats **trich** in
both women
and men¹



Ask your healthcare provider
if SOLOSEC® is right for you.

One day.
One dose.
That's it!



solosec
secnidazole
2g Oral Granules

For more information about SOLOSEC®, visit [Solosec.com](https://www.solosec.com)

Select Important Safety Information

- Before taking SOLOSEC®, tell your healthcare provider about all of your medical conditions, including if you
 - Are pregnant or plan to become pregnant.
 - Are breastfeeding or plan to breastfeed. Breastfeeding is not recommended. You should not breastfeed for 96 hours (4 days) after taking SOLOSEC®.

Please see complete Important Safety Information on page 11 of this brochure.

LUPIN

© 2022 Lupin Pharmaceuticals, Inc. All Rights Reserved.
Solosec® is a registered trademark owned by Lupin, Inc. PP-SOL-0677